arXiv:1509.04832v3 [math.AG] 16 Mar 2016

ON THE CANONICAL DEGREES OF GORENSTEIN THREEFOLDS OF GENERAL TYPE

RONG DU† AND YUN GAO††

Abstract. Let X be a Gorenstein minimal projective 3-fold with at worst locally factorial terminal singularities. Suppose that the canonical map is generically finite onto its image. C. Hacon showed that the canonical degree is universally bounded by 576. We improved Hacon's universal bound to 360. Moreover, we gave all the possible canonical degrees of X if X is an abelian cover over $\mathbb{P}^3$ and constructed all the examples with these canonical degrees.

1. Introduction

The study of the canonical maps of projective varieties of general type is one of the central problems in algebraic geometry. For the case of surfaces, Persson ([Per]) constructed a surface of general type with canonical degree 16 in 1978. About the same time, Beauville ([Bea]) proved that the degree of the canonical map is less than or equal to 36 and with the equality holds if and only if X is a ball quotient surface with $K_X^2 = 36$, $p_g = 3$, $q = 0$, and $|K_X|$ is base point free. Later, Xiao also found some restrictions on surfaces with high canonical degrees ([Xiao]). Since the canonical degree is bounded above, next interesting question is to determine which positive integers d's occur as the degrees of the canonical map. There are plenty of examples (see [Bea], [Cat1], [V-Z]) with canonical degrees being 2. For $d = 3$ and $d = 5$, Tan ([Tan]) and Pardini ([Par1]) constructed several surfaces independently. When $p_g(\Sigma) = 0$, Beauville ([Bea]) constructed surfaces with $\chi(\mathcal{O}_X)$ arbitrarily large and the canonical degrees $2, 4, 6$ and 8 . For $d = 9$, Tan also constructed a surface in

---

† The Research Sponsored by the National Natural Science Foundation of China (Grant No. 11471116), Science and Technology Commission of Shanghai Municipality (Grant No. 13dz2260400) and Shanghai Pujiang Program (Grant No. 12PJ1402400).

††The Research Sponsored by the National Natural Science Foundation of China (Grant No. 11271250,11271251) and SMC program of Shanghai Jiao Tong University.

Both authors are supported by China NSF (Grant No. 11531007).

[Tan]. Later, Casnati ([Cas])constructed surfaces of canonical degree from 3 to 9 as subvarieties in some projective bundle given by Pfaffans of alternating matrices. The authors ([D-G]) classified the surfaces whose canonical maps are abelian covers over $\mathbb{P}^2$ and constructed these surfaces by explicit defining equations. Recently, Rito in his series papers ([Rit],[Rit2],[Rit3]) constructed some new surfaces of general type with canonical degrees 12, 16 and 24 respectively.

In dimension at least three, the situation seems much less clear. M. Chen studied the canonical map of fiber type ([Ch],[C-H]) and posted an open problem in [Ch] as follows: Let X be a Gorenstein minimal projective 3-fold with at worst locally factorial terminal singularities. Suppose that the canonical map is generically finite onto its image. Is the generic degree of the canonical map universally upper bounded? Hacon gave a positive answer to Chen's problem. More precisely, he showed that the canonical degree is at most 576.

In this paper, we improve Hacon's upper bound by showing the following main theorem.

Theorem 1.1. *Let X be a Gorenstein minimal complex projective 3-fold of general type with locally factorial terminal singularities. Suppose that $|K_X|$ defines a generically finite map $\phi_X : X \dashrightarrow \mathbb{P}^{p_g-1}$, then deg $\phi_X \leq 360$ and with the equality holds if and only if $p_g(X) = 4$, $q(X) = 2$, $\chi(\omega_X) = 5$, $K^3 = 360$ and $|K_X|$ is base point free.*

Since the canonical degree is bounded above, it is quite interesting to consider a parallel problem as surfaces that which positive integers d's occur as the degrees of the canonical map of Gorenstein minimal projective 3-fold. As far as we know, there are quite few examples about 3-fold of general type with higher canonical degree. Cai ([Cai]) constructed some examples of 3-fold with canonical degrees 32 and 64 based on the existence of the surface with canonical degree 16 which was constructed by Persson. We show that if the canonical map is an abelian cover over $\mathbb{P}^3$ then the only possible canonical degrees of a Gorenstein minimal projective 3-fold are 2^m ($1 \leqslant m \leqslant 5$), by explicit constructions.

2. Proof of the main theorem

Let X be a Gorenstein minimal complex projective 3-fold of general type with locally factorial terminal singularities. Suppose that $|K_X|$ defines a generically finite map $\phi_X : X \dashrightarrow \mathbb{P}^{p_g-1}$. We will base on Hacon's beautiful arguments to improve the universal upper bound of the canonical degree.

Proof. Since ϕ_X is generically finite, one has that $p_g(X) \geq 4$. Let $d = deg\ \phi_X$. By the Miyaoka-Yau inequality ([Mi]), we have

$$d(p_g(X) - 3) \leq K_X^3 \leq 72\chi(\omega_X).$$

If we can show $\chi(\omega_X) \leq p_g(X) + 1$, then

$$d \leq 72\frac{\chi(\omega_X)}{p_g(X) - 3} \leq 72\frac{p_g(X) + 1}{p_g(X) - 3} = 72(1 + \frac{4}{p_g(X) - 3}) \leq 360. \quad (2.1)$$

If $q(X) \leq 2$, then $\chi(\omega_X) \leq p_g(X) + q(X) - 1 \leq p_g(X) + 1$.

Now we can assume hereafter that $q(X) \geq 3$. Consider the Albanese map alb_X of X and the Stein factorization f of alb_X as follows:

$$\begin{tikzcd} X \arrow[r, "f"] \arrow[dr, "alb_X"'] & Y \arrow[d] \\ & Alb(X) \end{tikzcd} .$$

By Hacon's argument (see the proof of [Ha], Theorem 1.1), one has

1. $\chi(\omega_X) \leq p_g(X)$, if $\dim Y \geq 2$;
2. $\chi(\omega_X) \leq p_g(X) + \chi(\omega_Y)$ and $\chi(\omega_Y)p_g(F) \leq p_g(X)$, where F is the general fiber of f, if $\dim Y = 1$.

Hence if $\dim Y \geq 2$, by (2.1), the statement holds. More precisely,

$$d \leq 72\frac{p_g(X)}{p_g(X) - 3} \leq 288.$$

We only need to consider $\dim Y = 1$.

If $p_g(F) \leq \dim X - 1$, then

$$h^0(\mathcal{O}_X(K_X) \otimes \mathcal{O}_F) \leq h^0(\mathcal{O}_X(K_X + F) \otimes \mathcal{O}_F) = h^0(\mathcal{O}_F(K_F)) \leq \dim X - 1,$$

which means that dim $\mathrm{Im}(\phi_X|_F) \leq \dim X - 2$, and hence dim $\mathrm{Im}\phi_X \leq \dim X - 1$, which contradicts the assumption that ϕ_X is generically finite. So we have that $p_g(F) \geq \dim X = 3$ and then $p_g(X) \geq \chi(\omega_Y)p_g(F) = (q(X) - 1)p_g(F) \geq 6$.

Therefore

$$d \leq 72\frac{\chi(\omega_X)}{p_g(X) - 3} \leq 72\frac{p_g(X) + \chi(\omega_Y)}{p_g(X) - 3} \leq 72(1 + \frac{1}{p_g(F)})\frac{p_g(X)}{p_g(X) - 3} \leq 192.$$

From the argument above, we know that the equality of (2.1) holds if and only if $p_g(X) = 4$, $q(X) = 2$, $\chi(\omega_X) = 5$, $K_X^3 = 360$ and $|K_X|$ is base point free. $\square$

Remark 2.1. If X is nonsingular and the canonical divisor K_X is ample then the equality in the main theorem holds if and only if X is a ball quotient. We guess that such a ball quotient with those invariants exists. For the parallel case of surfaces with the maximal canonical degree, the surface of general type with canonical degree 36 does exist which was constructed as some fake projective plane by S. Yeung recently ([Yeung]).

3. Canonical maps defined by abelain covers

The theory of cyclic covers of algebraic surfaces was studied first by Comessatti in [Com]. Then F. Catanese ([Cat2]) studied smooth abelian covers in the case $(\mathbb{Z}_2)^{\oplus 2}$ and R. Pardini ([Par2]) analyzed the general case. In this section, we shall recall some basic definitions and results for abelian covers and construct minimal 3-folds of general type whose canonical maps are abelian covers over $\mathbb{P}^3$. Since our point of view is to find the defining equations, we use the second author's expressions and notations appearing in [Gao].

Let $\varphi : X \to Y$ is an abelian cover associated to abelian group $G \cong \mathbb{Z}_{n_1} \oplus \cdots \oplus \mathbb{Z}_{n_k}$, i.e., function field $\mathbb{C}(X)$ of X is an abelian extension of the rational function field $\mathbb{C}(Y)$ with Galois group G. Without lose of generality, we can assume $n_1|n_2 \cdots |n_k$.

Definition 3.1. The dates of abelian cover over Y with group G are k effective divisors $D_1, \cdots, D_k$ and k linear equivalent relations

$$D_1 \sim n_1 L_1, \cdots, D_k \sim n_k L_k.$$

Let $\mathscr{L}_i = \mathscr{O}_Y(L_i)$ and f_i be the defining equation of D_i, i.e., $D_i = \mathrm{div}(f_i)$, where $f_i \in H^0(Y, \mathscr{L}_i^{n_i})$. Denote $\mathbf{V}(\mathscr{L}_i) = \mathbf{Spec} S(\mathscr{L}_i)$ to be the line bundle corresponding to $\mathscr{L}_i$, where $S(\mathscr{L}_i)$ is the sheaf of symmetric $\mathscr{O}_Y$ algebra. Let z_i be the fiber coordinate of $\mathbf{V}(\mathscr{L}_i)$. Then the abelian cover can be realized by the normalizing of surface V defined by the system of equations

$$z_1^{n_1} = f_1, \cdots, z_k^{n_k} = f_k.$$

So we have the following diagram:

$$\begin{tikzcd}
X \arrow[r, "\text{normalization}"] \arrow[drr, "\varphi"'] & V \arrow[r, hook] \arrow[dr] & \oplus_{i=1}^{k} \mathbf{V}(\mathscr{L}_i) \arrow[d, "p"] \\
& & Y.
\end{tikzcd}$$

Sometimes we call X is defined by these equations if there is no confusions in the context.

We summerize our main results as follows.

Theorem 3.2. *(See* [Gao]*) Denote by* $[Z]$ *the integral part of a* $\mathbb{Q}$*-divisor* Z*,* $-L_g = -\sum\limits_{i=1}^{k} g_i L_i + \left[\sum\limits_{i=1}^{k} \frac{g_i}{n_i} D_i\right]$*. Then*

$$\varphi_* \mathcal{O}_X \ = \bigoplus_{g \in G} \mathcal{O}_Y(-L_g). \tag{3.1}$$

where $g = (g_1, \cdots, g_k) \in G$*.*

So the decomposition of $\varphi_* \mathcal{O}_X$ is totally determined by the abelian cover.

Corollary 3.3. *If* X *is non-singular,* D *is the divisor on* Y*, then*

$$h^i(X, \varphi^* \mathcal{O}_Y(D)) = \sum_{g \in G} h^i(Y, \mathcal{O}_Y(D - L_g))$$

If the canonical map of X is an abelian cover over $\mathbb{P}^3$ then we have the explicit decomposition of $\varphi_* \mathcal{O}_X$.

Lemma 3.4. *If* $\varphi = \phi_X$ *is a finite abelian cover of degree* d *over* $\mathbb{P}^3$*, then* $\varphi_* \mathcal{O}_X = \mathcal{O}_{\mathbb{P}^3} \oplus \mathcal{O}_{\mathbb{P}^3}(-2)^{\oplus d/2-1} \oplus \mathcal{O}_{\mathbb{P}^3}(-3)^{\oplus d/2-1} \oplus \mathcal{O}_{\mathbb{P}^3}(-5)$*.*

Proof. Because φ is a finite abelian cover, $\varphi_* \mathcal{O}_X$ is a direct sum of the line bundles by Theorem 3.2.

$$\varphi_* \mathcal{O}_X = \mathcal{O}_{\mathbb{P}^3} \oplus \bigoplus_{i=1}^{d-1} \mathcal{O}_{\mathbb{P}^3}(-l_i).$$

Assume $0 < l_{d-1} \leqslant l_{d-2} \leqslant \cdots \leqslant l_1$.

Since $K_X = \varphi^*(\mathcal{O}_{\mathbb{P}^3}(1))$, for any $m \geqslant 1$,

$$P_m(X) = h^0(mK_X) = h^0(\varphi^*(\mathcal{O}_{\mathbb{P}^3}(m))) = h^0(\mathcal{O}_{\mathbb{P}^3}(m)) + \sum_{i=1}^{d-1} h^0(\mathcal{O}_{\mathbb{P}^3}(m - l_i)).$$

Because $p_g(X) = h^0(\varphi^*(\mathcal{O}_{\mathbb{P}^3}(H))) = h^0(\mathcal{O}_{\mathbb{P}^3}(1)) = 4$, we see that

$$h^0(\mathcal{O}_{\mathbb{P}^3}(1 - l_i)) = 0, \quad 1 \leq i \leq d - 1.$$

So $l_i \geqslant 2$.

And $p_g = h^3(\varphi_* \mathcal{O}_X) = h^3(\mathcal{O}_{\mathbb{P}^3}) + \sum_{i=1}^{d-1} h^3(\mathcal{O}_{\mathbb{P}^3}(-l_i))$,

So $4 = \sum_{i=1}^{d-1} h^0(\mathcal{O}_{\mathbb{P}^3}(l_i - 4))$, then $l_i \leqslant 5$.

Therefore, we have two cases as follows:

(1) $l_1 = 5, \quad l_2, \cdots, l_{d-1} < 4$, and

(2) $l_1 = l_2 = l_3 = l_4 = 4, \quad l_5, \cdots, l_{d-1} < 4$.

Let $m=2$, we have the second plurigenus of X $P_2(X)=\chi(2K_X)=\frac{1}{2}K_X^3+\frac{1}{6}K_Xc_2+\chi(\mathcal{O}_X)=\frac{d}{2}+3\chi(K_X)=\frac{d}{2}+9$. On the other hand, $P_2(X)=h^0(\mathcal{O}_{\mathbb{P}^3}(2))+\sum_{i=1}^{d-1}h^0\big(\mathcal{O}_{\mathbb{P}^3}(2-l_i)\big)$. So

$$\sum_{i=1}^{d-1}h^0(\mathcal{O}_{\mathbb{P}^3}(2-l_i))=\frac{d}{2}-1,$$

then there are exact $(\frac{d}{2}-1)$ 2's among l_i's.

Let $m=3$, we have the third plurigenus of X $P_3(X)=\chi(3K_X)=\frac{5}{2}K_X^3+\frac{1}{4}K_Xc_2+\chi(\mathcal{O}_X)=\frac{5d}{2}+5\chi(K_X)=\frac{5d}{2}+15$. On the other hand, $P_3(X)=h^0(\mathcal{O}_{\mathbb{P}^3}(3))+\sum_{i=1}^{d-1}h^0\big(\mathcal{O}_{\mathbb{P}^3}(3-l_i)\big)$. So

$$\sum_{i=1}^{d-1}h^0(\mathcal{O}_{\mathbb{P}^3}(3-l_i))=\frac{5d}{2}-5,$$

The second case does not satisfy the equation. So the lemma is proved. □

Now let $\varphi: X\to\mathbb{P}^3$ be an abelian cover associated to an abelian group $G\cong\mathbb{Z}_{n_1}\oplus\cdots\oplus\mathbb{Z}_{n_k}$. Then X is the normalization of the 3-fold defined by

$$z_1^{n_1}=f_1=\prod_\alpha p_\alpha^{\alpha_1},\cdots,z_k^{n_k}=f_k=\prod_\alpha p_\alpha^{\alpha_k},$$

where p_α's are coprime and $\alpha=(\alpha_1,\cdots,\alpha_k)\in G$. Denote x_α to be the degree of p_α, $e_i=(0,\cdots,0,1,0,\cdots,0)\in G$, $1\le i\le k$, and l_g be the degree of L_g, $g\in G$. So x_α and l_g are all integers. Then

$$n_il_{e_i}=\sum_\alpha\alpha_ix_\alpha\quad i=1,\cdots k,\tag{3.2}$$

$$l_g=\sum_{i=1}^k g_il_{e_i}-\sum_\alpha\left[\sum_{i=1}^k\frac{g_i\alpha_i}{n_i}\right]x_\alpha.\tag{3.3}$$

Lemma 3.5. *Using the notation as above, if $\varphi=\phi_X$, then there exists $g'=(g'_1,\cdots,g'_k)\in G\cong\mathbb{Z}_{n_1}\oplus\cdots\oplus\mathbb{Z}_{n_k}$ and a partition of G set-theoradically, $G=\{0\}\cup\{g'\}\cup S_1\cup S_2$, where the cardinalities of S_1*

and S_2 are equal, such that x_α satisfies the following equation

$$(*)\qquad \begin{cases} n_i l_{e_i} = \sum\limits_{\alpha} \alpha_i x_\alpha \\ l_{g'} = \sum\limits_{i=1}^{k} g'_i l_{e_i} - \sum\limits_{\alpha} \left[\sum\limits_{i=1}^{k} \dfrac{g'_i \alpha_i}{n_i} \right] x_\alpha = 5 \\ l_g = \sum\limits_{i=1}^{k} g_i l_{e_i} - \sum\limits_{\alpha} \left[\sum\limits_{i=1}^{k} \dfrac{g_i \alpha_i}{n_i} \right] x_\alpha = 3, \qquad g \in S_1 \\ l_g = \sum\limits_{i=1}^{k} g_i l_{e_i} - \sum\limits_{\alpha} \left[\sum\limits_{i=1}^{k} \dfrac{g_i \alpha_i}{n_i} \right] x_\alpha = 2, \qquad g \in S_2 \end{cases}$$

Proof. It comes from Lemma 3.4 directly.

□

By the above lemma, finding 3-folds whose canonical map are abelian covers over $\mathbb{P}^3$ is equivalent to finding the integral roots $\{x_\alpha\}$ of the above equations.

Theorem 3.6. *Let X be a Gorenstein minimal complex projective 3-fold of general type with locally factorial terminal singularities and $\varphi : X \longrightarrow \mathbb{P}^3$ is an abelian cover. If $\varphi = \phi_X$ then the canonical degree can only be 2^m, $1 \leqslant m \leqslant 5$.*

Proof. By Lemma 3.5, we only need to find the integral solutions of the equations (*). So the only possible degrees are $2, 4, 8, 16$ and 32 by using computer calculations. Moreover, we have the defining equations of the examples of all the degrees as follows.

Degree 2:

$$z^2 = f;$$

Degree 4:

$$\begin{cases} z_1^2 = s \\ z_2^2 = q; \end{cases}$$

Degree 8:

$$\begin{cases} z_1^2 = t_1 q \\ z_2^2 = t_2 q \\ z_3^2 = t_3 q; \end{cases}$$

Degree 16:

$$\begin{cases} z_1^2 = h_1 h_4 t_1 t_2 \\ z_2^2 = h_2 h_4 t_2 t_3 \\ z_3^2 = h_3 h_4 t_1 t_3 \\ z_4^2 = h_2 h_3 t_3; \end{cases}$$

Degree 32:

$$\begin{cases} z_1^2 = h_1h_2h_3h_{10} \\ z_2^2 = h_4h_5h_6h_{10} \\ z_3^2 = h_2h_3h_6h_7 \\ z_4^2 = h_1h_3h_5h_8 \\ z_5^2 = h_7h_8h_9h_{10}; \end{cases}$$

where the degree of h's is 1, t's is 2, q's is 4, s's is 6, f's is 10 and they all define nonsingular surfaces in $\mathbb{P}^3$ and intersect normal crossingly.

We want to show these 3-folds are smooth after normalization. Since the arguments are similar, we only prove the most complicated case with canonical degree 32.

Actually, we only need to consider the intersections of the branch locus locally. Let ℓ_{ij}'s be the intersection lines of h_i and h_j. Around the general point of ℓ_{ij} (except the intersection of three planes), the cover is locally defined by

$$z_1^2 = x^{a_{11}}y^{a_{12}}, \quad z_2^2 = x^{a_{21}}y^{a_{22}}, \quad z_3^2 = x^{a_{31}}y^{a_{32}}, \quad z_4^2 = x^{a_{41}}y^{a_{42}},$$

where $a_{ij} = 0$ or 1 for all i, j.

It is easy to check that $\{(a_{i1}, a_{i2})\} \subsetneqq \{(1,1),(0,0)\}$ i.e., at least one pair $\{(a_{i1}, a_{i2})\} = \{(1,0)\}$ or $\{(0,1)\}$. Without lose of generality, we can assume $(a_{11}, a_{21}) = (1,0)$, i.e. $z_1^2 = x$. After normalization, the cover is branched along the smooth surfaces. So the 3-fold is smooth at the preimages of the general points of ℓ_{ij}'s under the normalization map.

Let p_{ijk} be the intersection point of h_i, h_j and h_k. The arguments are similar. Let us take p_{123} for example. The cover is locally defined by

$$z_1^2 = xyw, \quad z_3^2 = yw, \quad z_4^2 = w.$$

After normalization, the cover is locally defined by

$$z_1^2 = x, \quad z_3^2 = y, \quad z_4^2 = w.$$

So the 3-folds are smooth.

It is easy to see that these 3-folds are all nonsingular with $p_g(X) = 4$, $q(X) = h^{2,0} = 0$, $\chi(\mathscr{O}_X) = -3$, $K_X = \varphi^*(\mathscr{O}_{\mathbb{P}^3}(1))$ and K_X^3 equals each degree of the covers. □

Remark 3.7. By Lemma 3.5, we have the integral solution of the equations (*) for degree 6 and 18. But the isolated singularities of corresponding 3-fold are not Gorenstein terminal since they are not cDV ([Rei2]).

Acknowledgements

Both authors would like to thank N. Mok for supporting their researches when they were in the University of Hong Kong and the referees for the useful comments.

References

[Bea] Beauville, A.: *L'application canonique pour les surfaces de type général*, Invention Math. **55**, 121-140(1979).

[Cai] Cai, J.: Degree of the canonical map of a Georenstein 3-fold of general type, Proc. AMS, **136**, (2008), No. 5, 1565-1574.

[Cas] Casnati, G.: *Covers of algebraic varieties. II. Covers of degree 5 and construction of surfaces.*, J. Algebraic Geom. **5** (1996), no. 3, 461-477.

[Cat1] Catanese, F.: *Babbage's conjecture, contact of surfaces, symmetric determinatal varieties and applications*, Invent. Math. **63** (1981), 433-465.

[Cat2] Catanese, F.: *On the moduli spaces of surfaces of general type*, J. Differential Geom., **19** (1984), 483-515.

[Ch] Chen, M.: *Weak Boundedness Theorems for Canonically Fibered Gorenstein Minimal 3-Folds*, Proc. Amer. Math. Soc., **133** (2005), No. 5, 1291-1298.

[C-H] Chen, M.; Hacon, C.: *On the geography of Gorenstein minimal 3-folds of general type*, Asian J. Math. **10** (2006), No.4, 757-764.

[Com] Comessatti, A.: *Sulle superfici multiple cicliche*, Rend. Sem. Mat. Univ. Padova, **1** (1930), 1-45.

[D-G] Du, R.; Gao, Y.: *Canonical maps of surfaces defined by abelian covers*, Asian J. Math. **18** (2014), No. 1, 219-228.

[Gao] Gao, Y.: *A note on finite abelian cover*, Science in China, Series A: Mathematics **54** (2011), 1333-1342.

[Ha] Hacon, C.: *On the degree of the canonical maps of 3-folds*, Proc. Japan Acad. **80** Ser. A (2004) 166-167.

[Mi] Miyaoka, Y.:*The Chern classes and Kodaira dimension of a minimal variety*, Algebraic geometry, Sendai, 1985, 449-476, Adv. Stud. Pure Math., **10**, North-Holland, Amsterdam, 1987.

[Par1] Pardini, R.: *Canonical images of surfaces*, J. reine angew. Math. **417**(1991), 215-219.

[Par2] Pardini, R.: *Abelian covers of algebraic varieties*, J. Reine Angew. Math. **417**(1991), 191-213.

[Per] Persson, U.: *Double coverings and surfaces of general type.*, In: Olson,L.D.(ed.) Algebraic geometry.(Lect. Notes Math., vol.732, pp.168-175) Berlin Heidelberg New York: Springer 1978.

[Rei] Reid, M.: *Surfaces with* $p_g = 0, K^2 = 2$. Preprint available at http://www.warwick.ac.uk/∼masda/surf/.

[Rei2] Reid, M., *Minimal models of canonical 3-folds*, Algebraic varieties and analytic varieties (Tokyo, 1981), 131C180, Adv. Stud. Pure Math., vol. 1, North-Holland, Amsterdam, 1983.

[Rit] Rito, C.: *New canonical triple covers of surfaces.* Proc. AMS, Vol. 143, No. 11 (2015), 4647-4653.

[Rit2] Rito, C.: *A surface with $q = 2$ and canonical map of degree* 16, arXiv:1506.05987.

[Rit3] Rito, C.: *A surface with canonical map of degree* 24, arXiv:1509.04132.

[Tan] Tan, S.-L.: *Surfaces whose canonical maps are of odd degrees.* Math. Ann. **292** (1992), no. 1, 13-29.

[V-Z] van Der Geer, G., Zagier, D: *The Hilbert modular group for the field* $\mathbb{Q}[\sqrt{13}]$, Invent. Math. **42** (1977), 93-134.

[Xiao] Xiao, G.: *Algebraic surfaces with high canonical degree*, Math. Ann. **274**(1986), 473-483.

[Yeung] Yeung, S.-L.: *A surface of maximal canonical degree* arXiv:1510.06622

Department of Mathematics, Shanghai Key Laboratory of PMMP, East China Normal University, Rm. 312, Math. Bldg, No. 500, Dongchuan Road, Shanghai, 200241, P. R. China
E-mail address: `rdu@math.ecnu.edu.cn`

Department of Mathematics, Shanghai Jiao Tong University, Shanghai 200240, P. R. of China
E-mail address: `gaoyunmath@sjtu.edu.cn`